


In the claims:

1. (Currently Amended) A compound of formula (I)




(I)


or a salt, ester, amide or prodrug thereof;

where X is O, or S, S(O), S(O)<sub>2</sub> or NR<sup>6</sup> where R<sup>6</sup> is hydrogen or C<sub>1-6</sub>alkyl;

R<sup>5</sup> is a group of sub-formulae (i) or (ii)




(i)




(ii)


or a group of sub-formula (iii), (iv) or (v)



(iii)



(iv)



(v)

where R<sup>80</sup> is a substituent of at least 4 atoms comprising one or more of:

1) halo, C<sub>1-4</sub>alkyl, optionally substituted C<sub>1-6</sub> alkoxy, C<sub>1-4</sub>alkoxymethyl,

di(C<sub>1-4</sub>alkoxy)methyl, C<sub>1-4</sub>alkanoyl, carboxy, benzoyl, trifluoromethyl, cyano, amino,


C<sub>2-5</sub>alkenyl, C<sub>2-5</sub>alkynyl, a phenyl group, a benzyl group or a 5-6-membered heterocyclic

group with 1-3 heteroatoms, selected independently from O, S and N, which heterocyclic

group may be aromatic or non-aromatic and may be saturated (linked via a ring carbon or

nitrogen atom) or unsaturated (linked via a ring carbon atom), and which phenyl, benzyl or heterocyclic group may bear on one or more ring carbon atoms up to 5 substituents selected from hydroxy, halogeno, C<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkoxy, C<sub>1-3</sub>alkanoyloxy, trifluoromethyl, cyano, amino, nitro, C<sub>2-4</sub>alkanoyl, C<sub>1-4</sub>alkanoylamino, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>alkylsulphonyl, C<sub>1-4</sub>alkylsulphinyl, C<sub>1-4</sub>alkylsulphonyl, carbamoyl, N-C<sub>1-4</sub>alkylcarbamoyl, N,N-di(C<sub>1-4</sub>alkyl)carbamoyl, aminosulphonyl, N-C<sub>1-4</sub>alkylaminosulphonyl, N,N-di(C<sub>1-4</sub>alkyl)aminosulphonyl, C<sub>1-4</sub>alkylsulphonylamino, and a saturated heterocyclic group selected from morpholino, thiomorpholino, pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl and pyrazolidinyl, which saturated heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, C<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkoxy, C<sub>1-3</sub>alkanoyloxy, trifluoromethyl, cyano, amino, nitro and C<sub>1-4</sub>alkoxycarbonyl;

2) a group of sub-formula (II)



where q' is 0, 1, 2, 3 or 4;

s' is 0 or 1;

X<sup>12</sup> is C(O) or S(O<sub>2</sub>),

R<sup>70</sup> is hydrogen, hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, amino, N-C<sub>1-6</sub>alkylamino,

N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>amino, hydroxyC<sub>2-6</sub>alkoxy, C<sub>1-6</sub>alkoxyC<sub>2-6</sub>alkoxy, aminoC<sub>2-6</sub>alkoxy,

N-C<sub>1-6</sub>alkylaminoC<sub>2-6</sub>alkoxy, N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>aminoC<sub>2-6</sub>alkoxy or C<sub>3-7</sub>cycloalkyl,

or R<sup>70</sup> is of the Formula (III):



wherein J is aryl, heteroaryl or heterocyclyl and K is a bond, oxy, imino,


N-(C<sub>1-6</sub>alkyl)imino, oxyC<sub>1-6</sub>alkylene, iminoC<sub>1-6</sub>alkylene, N-(C<sub>1-6</sub>alkyl)iminoC<sub>1-6</sub>alkylene,

-NHC(O)-, -SO<sub>2</sub>NH-, -NHSO<sub>2</sub>- or -NHC(O)-C<sub>1-6</sub>alkylene-,


and any aryl, heteroaryl or heterocyclyl group in a R<sup>70</sup> group may be optionally

substituted by one or more groups selected from hydroxy, oxo, halo, trifluoromethyl,

cyano, mercapto, nitro, amino, carboxy, carbamoyl, formyl, sulphamoyl, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>1-6</sub>alkoxy, -O-(C<sub>1-3</sub>alkyl)-O-, C<sub>1-6</sub>alkylS(O)<sub>n</sub>- (wherein n is 0-2), N-C<sub>1-6</sub>alkylamino, N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>amino, C<sub>1-6</sub>alkoxycarbonyl, N-C<sub>1-6</sub>alkylcarbamoyl, N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>2-6</sub>alkanoyl, C<sub>1-6</sub>alkanoyloxy, C<sub>1-6</sub>alkanoylamino, N-C<sub>1-6</sub>alkylsulphamoyl, N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-6</sub>alkylsulphonylamino and C<sub>1-6</sub>alkylsulphonyl-N-(C<sub>1-6</sub>alkyl)amino, or any aryl, heteroaryl or heterocyclyl group in a R<sup>70</sup> group may be optionally substituted with one or more groups of the Formula (IV):



wherein A<sup>1</sup> is halo, hydroxy, C<sub>1-6</sub>alkoxy, cyano, amino, N-C<sub>1-6</sub>alkylamino, N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>amino, carboxy, C<sub>1-6</sub>alkoxycarbonyl, carbamoyl, N-C<sub>1-6</sub>alkylcarbamoyl or N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>carbamoyl, p is 1 - 6, and B<sup>1</sup> is a bond, oxy, imino, N-(C<sub>1-6</sub>alkyl)imino or -NHC(O)-, with the proviso that p is 2 or more unless B<sup>1</sup> is a bond or -NHC(O)-; or any aryl, heteroaryl or heterocyclyl group in a R<sup>70</sup> group may be optionally substituted with one or more groups of the Formula (V):





wherein D<sup>1</sup> is aryl, heteroaryl or heterocyclyl and E<sup>1</sup> is a bond, C<sub>1-6</sub>alkylene, oxyC<sub>1-6</sub>alkylene, oxy, imino, N-(C<sub>1-6</sub>alkyl)imino, iminoC<sub>1-6</sub>alkylene, N-(C<sub>1-6</sub>alkyl)-iminoC<sub>1-6</sub>alkylene, C<sub>1-6</sub>alkylene-oxyC<sub>1-6</sub>alkylene, C<sub>1-6</sub>alkylene-iminoC<sub>1-6</sub>alkylene, C<sub>1-6</sub>alkylene-N-(C<sub>1-6</sub>alkyl)-iminoC<sub>1-6</sub>alkylene, -NHC(O)-, -NHSO<sub>2</sub>-, -SO<sub>2</sub>NH- or -NHC(O)-C<sub>1-6</sub>alkylene-, and any aryl, heteroaryl or heterocyclyl group in a substituent on R<sup>5</sup> may be optionally substituted with one or more groups selected from hydroxy, halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, carboxy, C<sub>1-6</sub>alkoxycarbonyl, carbamoyl, N-C<sub>1-6</sub>alkylcarbamoyl, N-(C<sub>1-6</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>2-6</sub>alkanoyl, amino, N-C<sub>1-6</sub>alkylamino and N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>amino, and any C<sub>3-7</sub>cycloalkyl or heterocyclyl group in a R<sup>70</sup> group may be optionally substituted with one or two oxo or thioxo substituents, and any of the R<sup>70</sup> groups defined hereinbefore which comprises a CH<sub>2</sub> group which is attached to 2 carbon atoms or a CH<sub>3</sub> group which is attached to a carbon atom may

optionally bear on each said CH<sub>2</sub> or CH<sub>3</sub> group a substituent selected from hydroxy, amino, C<sub>1-6</sub>alkoxy, N-C<sub>1-6</sub>alkylamino, N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>amino and heterocyclyl;

or R<sup>70</sup> may be cycloalkenyl or alkenyl optionally substituted by aryl;

and R<sup>99</sup> is hydrogen or a group C(O)R<sup>70</sup> where R<sup>70</sup> is as defined above;

3) a group of sub-formula (d) or (e)



where p' is 1-3, X<sup>10</sup> and X<sup>11</sup> are independently selected from a bond, -O-, -S- or NR<sup>101</sup>-


where R<sup>101</sup> is hydrogen or a C<sub>1-3</sub>alkyl, provided that one of X<sup>10</sup> or X<sup>11</sup> is a bond; X<sup>13</sup> is

-O-, -S- or NR<sup>102</sup>- where R<sup>102</sup> is hydrogen or a C<sub>1-3</sub>alkyl and R<sup>100</sup> is hydrogen or

optionally substituted hydrocarbyl or optionally substituted heterocyclyl, wherein any

optional substituents may be functional groups;

4) a group of formula (VI)



(VI)

where R<sup>71</sup> and R<sup>72</sup> are independently selected from hydrogen or C<sub>1-4</sub>alkyl, or R<sup>71</sup> and

R<sup>72</sup> together form a bond, and R<sup>73</sup> is a group OR<sup>74</sup>, NR<sup>75</sup>R<sup>76</sup> where R<sup>74</sup>, R<sup>75</sup> and R<sup>76</sup> are

independently selected from optionally substituted hydrocarbyl or optionally substituted

heterocyclic groups, and R<sup>75</sup> and R<sup>76</sup> may additionally form together with the nitrogen


atom to which they are attached, an aromatic or non-aromatic heterocyclic ring which

may contain further heteroatoms, wherein suitable optional substituents for hydrocarbyl

or heterocyclic groups R<sup>74</sup>, R<sup>75</sup> and R<sup>76</sup> include functional groups and heterocyclic groups

R<sup>74</sup>, R<sup>75</sup> and R<sup>76</sup> may further be substituted by a hydrocarbyl group;

5) a group of sub-formula (f)



(f)

where p'' is 0 or 1 and R<sup>83</sup> and R<sup>84</sup> are independently selected from hydrogen, optionally substituted hydrocarbyl or optionally substituted heterocyclyl, or R<sup>83</sup> and R<sup>84</sup> together with the nitrogen atom to which they are attached form an optionally substituted heterocyclic ring and where optional substituents for hydrocarbyl or heterocyclic groups R<sup>83</sup> and R<sup>84</sup> include functional groups and heterocyclic groups R<sup>83</sup> or R<sup>84</sup> may further be substituted by a hydrocarbyl group; and

R<sup>81</sup> is hydrogen, halo, C<sub>1-4</sub>alkoxy, cyano or trifluoromethyl, or phenyl an optionally substituted 6-membered aromatic ring containing at least one nitrogen atom, and R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> are independently selected from halogeno, cyano, nitro, C<sub>1-3</sub>alkylsulphanyl, -N(OH)R<sup>7</sup> (wherein R<sup>7</sup> is hydrogen, or C<sub>1-3</sub>alkyl), or R<sup>9</sup>X<sup>1</sup> (wherein X<sup>1</sup> represents a direct bond, -O-, -CH<sub>2</sub>-, -OC(O)-, -C(O)-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>10</sup>C(O)-, -C(O)NR<sup>11</sup>-, -SO<sub>2</sub>NR<sup>12</sup>-, -NR<sup>13</sup>SO<sub>2</sub>- or -NR<sup>14</sup> (wherein R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl)), and R<sup>9</sup> is hydrogen, optionally substituted hydrocarbyl, optionally substituted heterocyclyl or optionally substituted alkoxy where the optional substituents comprise at least one functional group; provided that at least one of R<sup>2</sup> or R<sup>3</sup> is other than hydrogen; and where a functional group is selected from nitro, cyano, halo, oxo, =CR<sup>78</sup>R<sup>79</sup>, C(O)<sub>x</sub>R<sup>77</sup>, OR<sup>77</sup>, S(O)<sub>y</sub>R<sup>77</sup>, NR<sup>78</sup>R<sup>79</sup>, C(O)NR<sup>78</sup>R<sup>79</sup>, OC(O)NR<sup>78</sup>R<sup>79</sup>, =NOR<sup>77</sup>, -NR<sup>77</sup>C(O)<sub>x</sub>R<sup>78</sup>, -NR<sup>77</sup>CONR<sup>78</sup>R<sup>79</sup>, -N=CR<sup>78</sup>R<sup>79</sup>, S(O)<sub>y</sub>NR<sup>78</sup>R<sup>79</sup> or -NR<sup>77</sup>S(O)<sub>y</sub>R<sup>78</sup> where R<sup>77</sup>, R<sup>78</sup> and R<sup>79</sup> are independently selected from hydrogen, optionally substituted hydrocarbyl, optionally substituted heterocyclyl or optionally substituted alkoxy, or R<sup>78</sup> and R<sup>79</sup> together form an optionally substituted ring which optionally contains further heteroatoms such as oxygen, nitrogen, S, S(O) or S(O)<sub>2</sub>, where x is an integer of 1 or 2, y is 0 or an integer of 1-3 and where hydrocarbyl, heterocyclyl or alkoxy groups R<sup>77</sup>, R<sup>78</sup> and R<sup>79</sup> as well as rings formed by R<sup>78</sup> and R<sup>79</sup> are optionally substituted by halo, perhaloalkyl, mercapto, alkylthio, hydroxy, carboxy, alkoxy, heteroaryl, heteroaryloxy,

cycloalkyl, cycloalkenyl, cycloalkynyl, alkenyloxy, alkynyloxy, alkoxyalkoxy, aryloxy (where the aryl group may be substituted by halo, nitro, or hydroxy), cyano, nitro, amino, mono- or di-alkyl amino, oximino or S(O)<sub>y</sub>R<sup>90</sup> where y is 0 or an integer of 1-3 and R<sup>90</sup> is a alkyl; and  
wherein hydrocarbyl is selected from alkyl, alkenyl, alkynyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, or combinations thereof.

2. (Canceled)
3. (Canceled)
4. (Currently Amended) A compound according to claim 1 ~~any one of the preceding claims~~ wherein at least one group R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> is a group R<sup>9</sup>X<sup>1</sup>- and R<sup>9</sup> is hydrogen or an alkyl group, optionally substituted with one or more groups selected from functional groups as defined in claim ~~12 or claim 3~~, or alkenyl, alkynyl, aryl, heterocyclyl, cycloalkyl, or cycloalkenyl or cycloalkynyl, any of which may be substituted with a functional group as defined in claim ~~12 or claim 3~~, and where any aryl, heterocyclyl, cycloalkyl, or cycloalkenyl, cycloalkynyl groups may also be optionally substituted with ~~hydrocarbyl such as~~ alkyl, alkenyl or alkynyl.
5. (Currently Amended) A compound according to claim 1 wherein at least one group R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> is a group R<sup>9</sup>X<sup>1</sup>- and R<sup>9</sup> is selected from one of the following twenty-two groups:
  - 1) hydrogen or C<sub>1-5</sub>alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, oxiranyl, fluoro, chloro, bromo and amino (including C<sub>1-3</sub>alkyl and trifluoromethyl);
  - 2) -R<sup>a</sup>X<sup>2</sup>C(O)R<sup>15</sup> (wherein X<sup>2</sup> represents -O- or -NR<sup>16</sup>- (in which R<sup>16</sup> represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>15</sup> represents C<sub>1-3</sub>alkyl, -NR<sup>17</sup>R<sup>18</sup> or -OR<sup>19</sup> (wherein R<sup>17</sup>, R<sup>18</sup> and R<sup>19</sup> which may be the same or different each represents hydrogen, C<sub>1-5</sub>alkyl, ~~hydroxyC<sub>1-5</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl~~));

- 3)  $-R^bX^3R^{20}$  (wherein  $X^3$  represents  $-O-$ ,  $C(O)-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-OC(O)-$ ,  $-NR^{21}C(O)_s-$ ,  $-C(O)NR^{22}-$ ,  $-SO_2NR^{23}-$ ,  $-NR^{24}SO_2-$  or  $-NR^{25}-$  (wherein  $R^{21}$ ,  $R^{22}$ ,  $R^{23}$ ,  $R^{24}$  and  $R^{25}$  each independently represents hydrogen,  $C_{1-3}$ alkyl, hydroxy  $C_{1-4}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl and  $s$  is 1 or 2) and  $R^{20}$  represents hydrogen,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which  $C_{1-6}$ alkyl group may bear 1, 2 or 3 substituents selected from oxo, hydroxy, halogeno, cyclopropyl, amino,  $C_{1-4}$ alkylamino,  $C_{1-4}$ alkanoyldi- $C_{1-4}$ alkylamino-,  $C_{1-4}$ alkylthio,  $C_{1-4}$ alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano,  $C_{1-4}$ cyanoalkyl,  $C_{1-4}$ alkyl,  $C_{1-4}$ hydroxyalkyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkoxy $C_{1-4}$ alkyl,  $C_{1-4}$ alkylsulphonyl $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxycarbonyl,  $C_{1-4}$ aminoalkyl,  $C_{1-4}$ alkylamino, di( $C_{1-4}$ alkyl)amino,  $C_{1-4}$ alkylamino $C_{1-4}$ alkyl, di( $C_{1-4}$ alkyl)amino $C_{1-4}$ alkyl,  $C_{1-4}$ alkylamino $C_{1-4}$ alkoxy, di( $C_{1-4}$ alkyl)amino $C_{1-4}$ alkoxy and a group  $-(-O-)_f(R^{b'})_gD$  (wherein  $f$  is 0 or 1,  $g$  is 0 or 1 and  $D$  is a  $C_{3-6}$ cycloalkyl group or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from  $C_{1-4}$ alkyl));
- 4)  $-R^cX^4R^cX^5R^{26}$  (wherein  $X^4$  and  $X^5$  which may be the same or different are each  $-O-$ ,  $C(O)-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^{27}C(O)_s-$ ,  $-C(O)_s[[X]]NR^{28}-$ ,  $-SO_2NR^{29}-$ ,  $-NR^{30}SO_2-$  or  $-NR^{31}-$  (wherein  $R^{27}$ ,  $R^{28}$ ,  $R^{29}$ ,  $R^{30}$  and  $R^{31}$  each independently represents hydrogen,  $C_{1-3}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl and  $s$  is 1 or 2) and  $R^{26}$  represents hydrogen,  $C_{1-3}$ alkyl, hydroxy $C_{1-3}$ alkyl\_or\_ $C_{1-3}$ alkoxy $C_{2-3}$ alkyl);
- 5)  $R^{32}$  (wherein  $R^{32}$  is a 4-6-membered cycloalkyl or saturated heterocyclic ring (linked via carbon or nitrogen) with 1-2 heteroatoms, selected independently from O, S and N, which cycloalkyl or heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano,  $C_{1-4}$ alkyl, hydroxy $C_{1-4}$ alkyl, cyano $C_{1-4}$ alkyl, cyclopropyl,  $C_{1-4}$ alkylsulphonyl $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxycarbonyl, carboxamido,  $C_{1-4}$ aminoalkyl,  $C_{1-4}$ alkylamino, di( $C_{1-4}$ alkyl)amino,  $C_{1-4}$ alkylamino $C_{1-4}$ alkyl,  $C_{1-4}$ alkanoyl, di( $C_{1-4}$ alkyl)amino $C_{1-4}$ alkyl,  $C_{1-4}$ alkylamino $C_{1-4}$ alkoxy, di( $C_{1-4}$ alkyl)amino $C_{1-4}$ alkoxy nitro, amino,  $C_{1-4}$ alkoxy,  $C_{1-4}$ hydroxyalkoxy, carboxy, trifluoromethyl,  $-C(O)NR^{38}R^{39}$ ,  $-NR^{40}C(O)R^{41}$  (wherein  $R^{38}$ ,  $R^{39}$ ,  $R^{40}$  and  $R^{41}$ , which may be the same or different, each represents hydrogen,  $C_{1-4}$ alkyl, hydroxy $C_{1-4}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl) and a group

$-(\text{-O-})_f(\text{C}_{1-4}\text{alkyl})_g\text{ringD}$  (wherein  $f$  is 0 or 1,  $g$  is 0 or 1 and ring D is a cyclic group selected from  $\text{C}_{3-6}$ cycloalkyl, aryl or 5-6-membered saturated or unsaturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from halo and  $\text{C}_{1-4}$ alkyl);

6)  $-\text{R}^d\text{R}^{32}$  (wherein  $\text{R}^{32}$  is as defined hereinbefore);

7)  $-\text{R}^e\text{R}^{32}$  (wherein  $\text{R}^{32}$  is as defined hereinbefore);

8)  $-\text{R}^f\text{R}^{32}$  (wherein  $\text{R}^{32}$  is as defined hereinbefore);

9)  $\text{R}^{33}$  (wherein  $\text{R}^{33}$  represents a pyridone group, a phenyl group or a 5-6-membered aromatic heterocyclic group (linked via carbon or nitrogen) with 1-3 heteroatoms selected from O, N and S, which pyridone, phenyl or aromatic heterocyclic group may carry up to 5 substituents selected from hydroxy, nitro, halogeno, amino,  $\text{C}_{1-4}$ alkyl,  $\text{C}_{1-4}$ alkoxy,  $\text{C}_{1-4}$ hydroxyalkyl,  $\text{C}_{1-4}$ aminoalkyl,  $\text{C}_{1-4}$ alkylamino,  $\text{C}_{1-4}$ hydroxyalkoxy, oxo, cyano $\text{C}_{1-4}$ alkyl, cyclopropyl,  $\text{C}_{1-4}$ alkylsulphonyl $\text{C}_{1-4}$ alkyl,  $\text{C}_{1-4}$ alkoxycarbonyl, di( $\text{C}_{1-4}$ alkyl)amino,  $\text{C}_{1-4}$ alkylamino $\text{C}_{1-4}$ alkyl,  $\text{C}_{1-4}$ alkanoyl, di( $\text{C}_{1-4}$ alkyl)amino $\text{C}_{1-4}$ alkyl,  $\text{C}_{1-4}$ alkylamino $\text{C}_{1-4}$ alkoxy, di( $\text{C}_{1-4}$ alkyl)amino $\text{C}_{1-4}$ alkoxy, carboxy, carboxamido, trifluoromethyl, cyano,  $-\text{C}(\text{O})\text{NR}^{38}\text{R}^{39}$ ,  $-\text{NR}^{40}\text{C}(\text{O})\text{R}^{41}$  (wherein  $\text{R}^{38}$ ,  $\text{R}^{39}$ ,  $\text{R}^{40}$  and  $\text{R}^{41}$ , which may be the same or different, each represents hydrogen,  $\text{C}_{1-4}$ alkyl, hydroxy $\text{C}_{1-4}$ alkyl or  $\text{C}_{1-3}$ alkoxy $\text{C}_{2-3}$ alkyl) and a group  $-(\text{-O-})_f(\text{C}_{1-4}\text{alkyl})_g\text{ringD}$  (wherein  $f$  is 0 or 1,  $g$  is 0 or 1 and ring D is a cyclic group selected from  $\text{C}_{3-6}$ cycloalkyl, aryl or 5-6-membered saturated or unsaturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from halo and  $\text{C}_{1-4}$ alkyl);

10)  $-\text{R}^g\text{R}^{33}$  (wherein  $\text{R}^{33}$  is as defined hereinbefore);

11)  $-\text{R}^h\text{R}^{33}$  (wherein  $\text{R}^{33}$  is as defined hereinbefore);

12)  $-\text{R}^i\text{R}^{33}$  (wherein  $\text{R}^{33}$  is as defined hereinbefore);

13)  $-\text{R}^j\text{X}^6\text{R}^{33}$  (wherein  $\text{X}^6$  represents  $-\text{O}-$ ,  $-\text{C}(\text{O})-$ ,  $-\text{S}-$ ,  $-\text{SO}-$ ,  $-\text{SO}_2-$ ,  $-\text{OC}(\text{O})-$ ,  $-\text{NR}^{38}\text{C}(\text{O})-$ ,  $-\text{C}(\text{O})\text{NR}^{39}-$ ,  $-\text{SO}_2\text{NR}^{40}-$ ,  $-\text{NR}^{41}\text{SO}_2-$  or  $-\text{NR}^{42}-$  (wherein  $\text{R}^{38}$ ,  $\text{R}^{39}$ ,  $\text{R}^{40}$ ,  $\text{R}^{41}$  and  $\text{R}^{42}$  each independently represents hydrogen,  $\text{C}_{1-3}$ alkyl, hydroxy $\text{C}_{1-3}$ alkyl or  $\text{C}_{1-3}$ alkoxy $\text{C}_{2-3}$ alkyl) and  $\text{R}^{33}$  is as defined hereinbefore);

14)  $-\text{R}^k\text{X}^7\text{R}^{33}$  (wherein  $\text{X}^7$  represents  $-\text{O}-$ ,  $\text{C}(\text{O})$ ,  $-\text{S}-$ ,  $-\text{SO}-$ ,  $-\text{SO}_2-$ ,  $-\text{NR}^{43}\text{C}(\text{O})-$ ,



- C(O)NR<sup>44</sup>-, -SO<sub>2</sub>NR<sup>45</sup>-, -NR<sup>46</sup>SO<sub>2</sub>- or -NR<sup>47</sup>- (wherein R<sup>43</sup>, R<sup>44</sup>, R<sup>45</sup>, R<sup>46</sup> and R<sup>47</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl, hydroxyC<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>33</sup> is as defined hereinbefore);
- 15) -R<sup>m</sup>X<sup>8</sup>R<sup>33</sup> (wherein X<sup>8</sup> represents -O-, -C(O)-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>48</sup>C(O)-, -C(O)NR<sup>49</sup>-, -SO<sub>2</sub>NR<sup>50</sup>-, -NR<sup>51</sup>SO<sub>2</sub>- or -NR<sup>52</sup>- (wherein R<sup>48</sup>, R<sup>49</sup>, R<sup>50</sup>, R<sup>51</sup> and R<sup>52</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl, hydroxyC<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>33</sup> is as defined hereinbefore);
- 16) -R<sup>n</sup>X<sup>9</sup>R<sup>n'</sup>R<sup>33</sup> (wherein X<sup>9</sup> represents -O-, -C(O)-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>53</sup>C(O)-, -C(O)NR<sup>54</sup>-, -SO<sub>2</sub>NR<sup>55</sup>-, -NR<sup>56</sup>SO<sub>2</sub>- or -NR<sup>57</sup>- (wherein R<sup>53</sup>, R<sup>54</sup>, R<sup>55</sup>, R<sup>56</sup> and R<sup>57</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl, hydroxyC<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>33</sup> is as defined hereinbefore);
- 17) -R<sup>p</sup>X<sup>9</sup>-R<sup>p'</sup>R<sup>32</sup> (wherein X<sup>9</sup> and R<sup>32</sup> are as defined hereinbefore);
- 18) C<sub>2-5</sub>alkenyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C<sub>1-4</sub>alkylamino, N,N-di(C<sub>1-4</sub>alkyl)amino, aminosulphonyl, N-C<sub>1-4</sub>alkylaminosulphonyl and N,N-di(C<sub>1-4</sub>alkyl)aminosulphonyl;
- 19) C<sub>2-5</sub>alkynyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C<sub>1-4</sub>alkylamino, N,N-di(C<sub>1-4</sub>alkyl)amino, aminosulphonyl, N-C<sub>1-4</sub>alkylaminosulphonyl and N,N-di(C<sub>1-4</sub>alkyl)aminosulphonyl;
- 20) -R<sup>t</sup>X<sup>9</sup>R<sup>t'</sup>R<sup>32</sup> (wherein X<sup>9</sup> and R<sup>32</sup> are as defined hereinbefore);
- 21) -R<sup>u</sup>X<sup>9</sup>R<sup>u'</sup>R<sup>32</sup> (wherein X<sup>9</sup> and R<sup>32</sup> are as defined hereinbefore); and
- 22) -R<sup>v</sup>R<sup>58</sup>(R<sup>v'</sup>)<sub>q</sub>(X<sup>9</sup>)<sub>r</sub>R<sup>59</sup> (wherein X<sup>9</sup> is as defined hereinbefore, q is 0 or 1, r is 0 or 1, and R<sup>58</sup> is a C<sub>1-3</sub>alkylene group or a cyclic group selected from cyclopropyl, cyclobutyl, cyclopentylene, cyclohexylene or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C<sub>1-3</sub>alkylene group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C<sub>1-4</sub>alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-4</sub>cyanoalkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylsulphonylC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>aminoalkyl, C<sub>1-4</sub>alkylamino, di(C<sub>1-4</sub>alkyl)amino, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkyl, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkyl,

$C_{1-4}$ alkylamino $C_{1-4}$ alkoxy, di( $C_{1-4}$ alkyl)amino $C_{1-4}$ alkoxy and a group  
 $-(O)_f(C_{1-4}alkyl)_g ringD$  (wherein  $f$  is 0 or 1,  $g$  is 0 or 1 and ring  $D$  is a cyclic group  
 selected from  $C_{3-6}$ cycloalkyl, aryl or 5-6-membered saturated or unsaturated heterocyclic  
 group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group  
 may bear one or more substituents selected from halo and  $C_{1-4}alkyl$ ); and  $R^{59}$  is hydrogen,  
 $C_{1-3}alkyl$ , or a cyclic group selected from cyclopropyl, cyclobutyl, cyclopentyl,  
 cyclohexyl and a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms,  
 selected independently from O, S and N, which  $C_{1-3}alkyl$  group may bear 1 or 2  
 substituents selected from oxo, hydroxy, halogeno,  $C_{1-4}alkoxy$  and which cyclic group  
 may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano,  
 $C_{1-4}cyanoalkyl$ ,  $C_{1-4}alkyl$ ,  $C_{1-4}hydroxyalkyl$ ,  $C_{1-4}alkoxy$ ,  $C_{1-4}alkoxyC_{1-4}alkyl$ ,  
 $C_{1-4}alkylsulphonylC_{1-4}alkyl$ ,  $C_{1-4}alkoxycarbonyl$ ,  $C_{1-4}aminoalkyl$ ,  $C_{1-4}alkylamino$ ,  
 di( $C_{1-4}alkyl$ )amino,  $C_{1-4}alkylaminoC_{1-4}alkyl$ , di( $C_{1-4}alkyl$ )amino $C_{1-4}alkyl$ ,  
 $C_{1-4}alkylaminoC_{1-4}alkoxy$ , di( $C_{1-4}alkyl$ )amino $C_{1-4}alkoxy$  and a group  
 $-(O)_f(C_{1-4}alkyl)_g ringD$  (wherein  $f$  is 0 or 1,  $g$  is 0 or 1 and  $D$  is a cyclic group selected  
 from  $C_{3-6}$ cycloalkyl, aryl or 5-6-membered saturated or unsaturated heterocyclic group  
 with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may  
 bear one or more substituents selected from halo and  $C_{1-4}alkyl$ );  
 and wherein  $R^a$ ,  $R^b$ ,  $R^{b'}$ ,  $R^c$ ,  $R^{c'}$ ,  $R^d$ ,  $R[[g]]_e$ ,  $R^j$ ,  $R^n$ ,  $R^{n'}$ ,  $R^p$ ,  $R^{p'}$ ,  $R^t$ ,  $R^u$ ,  $R^v$  and  $R^{v'}$  are  
 independently selected from  $C_{1-8}alkylene$  groups optionally substituted by one or more  
 substituents selected from hydroxy, halogeno, amino,  
 $R^e$ ,  $R^h$ ,  $R^k$  and  $R^l$  are independently selected from  $C_{2-8}alkenylene$  groups optionally  
 substituted by one or more substituents selected from hydroxy, halogeno, amino, and  
 $R^l$  may additionally be a bond; and  
 $R^f$ ,  $R^i$ ,  $R^m$  and  $R^u$  are independently selected from  $C_{2-8}alkynylene$  groups optionally  
 substituted by one or more substituents selected from hydroxy, halogeno, amino.

6. (Currently Amended) A compound according to claim 5 wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$  are independently selected from, halo, cyano, nitro, trifluoromethyl,  $C_{1-3}alkyl$ ,  $-NR^7R^8$  (wherein  $R^7$  and  $R^8$ , which may be the same or different, each represents hydrogen or  $C_{1-3}alkyl$ ), or other groups from formula  $-X^1R^9$  (wherein  $X^1$  represents a direct bond,  $-O-$ ,

-CH<sub>2</sub>-, -OCO-, carbonyl, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>10</sup>CO-, -CONR<sup>11</sup>-, -SO<sub>2</sub>NR<sup>12</sup>-, -NR<sup>13</sup>SO<sub>2</sub>- or -NR<sup>14</sup>- (wherein R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl), and R<sup>9</sup> is selected from one of the following groups:

1') hydrogen or C<sub>1-5</sub>alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro or amino,

2') C<sub>1-5</sub>alkylX<sup>2</sup>C(O)R<sup>15</sup> (wherein X<sup>2</sup> represents -O- or -NR<sup>16</sup>- (in which R<sup>15</sup> represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>16</sup> represents C<sub>1-3</sub>alkyl, -NR<sup>17</sup>R<sup>18</sup> or -OR<sup>19</sup> (wherein R<sup>17</sup>, R<sup>18</sup> and R<sup>19</sup> which may be the same or different each represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl));

3') C<sub>1-5</sub>alkylX<sup>3</sup>R<sup>20</sup> (wherein X<sup>3</sup> represents -O-, -S-, -SO-, -SO<sub>2</sub>-, -OCO-, -NR<sup>21</sup>CO-, -CONR<sup>22</sup>-, -SO<sub>2</sub>NR<sup>23</sup>-, -NR<sup>24</sup>SO<sub>2</sub>- or -NR<sup>25</sup>- (wherein R<sup>21</sup>, R<sup>22</sup>, R<sup>23</sup>, R<sup>24</sup> and R<sup>25</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>20</sup> represents hydrogen, C<sub>1-3</sub>alkyl, cyclopentyl, cyclohexyl or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C<sub>1-3</sub>alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C<sub>1-4</sub>alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl and C<sub>1-4</sub>alkoxy);

4') C<sub>1-5</sub>alkylX<sup>4</sup>C<sub>1-5</sub>alkylX<sup>5</sup>R<sup>26</sup> (wherein X<sup>4</sup> and X<sup>5</sup> which may be the same or different are each -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>27</sup>CO-, -CONR<sup>28</sup>-, -SO<sub>2</sub>NR<sup>29</sup>-, -NR<sup>30</sup>SO<sub>2</sub>- or -NR<sup>31</sup>- (wherein R<sup>27</sup>, R<sup>28</sup>, R<sup>29</sup>, R<sup>30</sup> and R<sup>31</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>26</sup> represents hydrogen or C<sub>1-3</sub>alkyl);

5') R<sup>32</sup> (wherein R<sup>32</sup> is a 5-6-membered saturated heterocyclic group (linked via carbon or nitrogen) with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl and C<sub>1-4</sub>alkylsulphonylC<sub>1-4</sub>alkyl);

6') C<sub>1-5</sub>alkylR<sup>32</sup> (wherein R<sup>32</sup> is as defined in (5') above);

7') C<sub>2-5</sub>alkenylR<sup>32</sup> (wherein R<sup>32</sup> is as defined in (5') above);

8') C<sub>2-5</sub>alkynylR<sup>32</sup> (wherein R<sup>32</sup> is as defined in (5') above);

9')  $R^{33}$  (wherein  $R^{33}$  represents a pyridone group, a phenyl group or a 5-6-membered aromatic heterocyclic group (linked via carbon or nitrogen) with 1-3 heteroatoms selected from O, N and S, which pyridone, phenyl or aromatic heterocyclic group may carry up to 5 substituents on an available carbon atom selected from hydroxy, halogeno, amino,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ hydroxyalkyl,  $C_{1-4}$ aminoalkyl,  $C_{1-4}$ alkylamino,  $C_{1-4}$ hydroxyalkoxy, carboxy, trifluoromethyl, cyano,  $-CONR^{34}R^{35}$  and  $-NR^{36}COR^{37}$  (wherein  $R^{34}$ ,  $R^{35}$ ,  $R^{36}$  and  $R^{37}$ , which may be the same or different, each represents hydrogen,  $C_{1-4}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl));

10')  $C_{1-5}$ alkyl $R^{33}$  (wherein  $R^{33}$  is as defined in (9') above);

11')  $C_{2-5}$ alkenyl $R^{33}$  (wherein  $R^{33}$  is as defined in (9') above);

12')  $C_{2-5}$ alkynyl $R^{33}$  (wherein  $R^{33}$  is as defined in (9') above);

13')  $C_{1-5}$ alkyl $X^6R^{33}$  (wherein  $X^6$  represents  $-O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^{38}CO-$ ,  $-CONR^{39}-$ ,  $-SO_2NR^{40}-$ ,  $-NR^{41}SO_2-$  or  $-NR^{42}-$  (wherein  $R^{38}$ ,  $R^{39}$ ,  $R^{40}$ ,  $R^{41}$  and  $R^{42}$  each independently represents hydrogen,  $C_{1-3}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl) and  $R^{33}$  is as defined hereinbefore);

14')  $C_{2-5}$ alkenyl $X^7R^{33}$  (wherein  $X^7$  represents  $-O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^{43}CO-$ ,  $-CONR^{44}-$ ,  $-SO_2NR^{45}-$ ,  $-NR^{46}SO_2-$  or  $-NR^{47}-$  (wherein  $R^{43}$ ,  $R^{44}$ ,  $R^{45}$ ,  $R^{46}$  and  $R^{47}$  each independently represents hydrogen,  $C_{1-3}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl) and  $R^{33}$  is as defined hereinbefore);


15')  $C_{2-5}$ alkynyl $X^8R^{33}$  (wherein  $X^8$  represents  $-O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^{48}CO-$ ,  $-C(O)NR^{49}-$ ,  $-SO_2NR^{50}-$ ,  $-NR^{51}SO_2-$  or  $-NR^{52}-$  (wherein  $R^{48}$ ,  $R^{49}$ ,  $R^{50}$ ,  $R^{51}$  and  $R^{52}$  each independently represents hydrogen,  $C_{1-3}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl) and  $R^{33}$  is as defined hereinbefore);

16')  $C_{1-3}$ alkyl $X^9C_{1-3}$ alkyl $R^{33}$  (wherein  $X^9$  represents  $-O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^{53}CO-$ ,  $-C(O)NR^{54}-$ ,  $-SO_2NR^{55}-$ ,  $-NR^{56}SO_2-$  or  $-NR^{57}-$  (wherein  $R^{53}$ ,  $R^{54}$ ,  $R^{55}$ ,  $R^{56}$  and  $R^{57}$  each independently represents hydrogen,  $C_{1-3}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl) and  $R^{33}$  is as defined hereinbefore); and

17')  $C_{1-3}$ alkyl $X^9C_{1-3}$ alkyl $R^{32}$  (wherein  $X^9$  and  $R^{32}$  are as defined in (5') above), provided that at least one of  $R^2$  or  $R^3$  is other than hydrogen.

7. (Currently Amended) A compound according to any one of ~~the preceding~~ claims 1, 4, 5 or 6, where  $R^1$  is hydrogen and  $R^4$  is hydrogen, halo,  $C_{1-4}$  alkyl or  $C_{1-4}$  alkoxy.
8. (Currently Amended) A compound according to any one of ~~the preceding~~ claims 1, 4, 5 or 6, wherein at least one group  $R^2$  or  $R^3$  comprises a chain of at least 3 optionally substituted carbon atoms or heteroatoms selected from oxygen, nitrogen or sulphur.
9. (Currently Amended) A compound according to claim 8 wherein said chain is substituted by a polar group ~~as defined in claim 5~~ which assists solubility, wherein the polar group is selected from oxo, hydroxy, halogeno, amino,  $C_{1-4}$  alkylamino,  $C_{1-4}$  alkanoyldi- $C_{1-4}$  alkylamino-,  $C_{1-4}$  alkylthio,  $C_{1-4}$  alkoxy, cyano,  $C_{1-4}$  cyanoalkyl,  $C_{1-4}$  hydroxyalkyl,  $C_{1-4}$  alkoxy $C_{1-4}$  alkyl,  $C_{1-4}$  alkylsulphonyl $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxycarbonyl,  $C_{1-4}$  aminoalkyl, di( $C_{1-4}$  alkyl)amino,  $C_{1-4}$  alkylamino $C_{1-4}$  alkyl, di( $C_{1-4}$  alkyl)amino $C_{1-4}$  alkyl,  $C_{1-4}$  alkylamino $C_{1-4}$  alkoxy, di( $C_{1-4}$  alkyl)amino $C_{1-4}$  alkoxy, nitro,  $C_{1-4}$  hydroxyalkoxy, di( $C_{1-4}$  alkyl)amino $C_{1-4}$  alkoxy, carboxy, carboxamido, trifluoromethyl,  $-C(O)NR^{38}R^{39}$ , and  $-NR^{40}C(O)R^{41}$  (wherein  $R^{38}$ ,  $R^{39}$ ,  $R^{40}$  and  $R^{41}$ , which may be the same or different, each represents hydrogen,  $C_{1-4}$  alkyl, hydroxy $C_{1-4}$  alkyl or  $C_{1-3}$  alkoxy $C_{2-3}$  alkyl).
10. (Currently Amended) A compound according to any one of ~~the preceding~~ claims 1, 4, 5 or 6, wherein  $R^3$  is a group  $X^1R^9$  where  $X^1$  is oxygen and  $R^9$  includes a methylene group directly adjacent  $X^1$ .
11. (Original) A compound according to claim 5 wherein at least one of  $R^1$ ,  $R^2$ ,  $R^3$  or  $R^4$  is a group  $X^1R^9$  which includes a bridging alkylene, alkenylene or alkynylene groups  $R^a$ ,  $R^b$ ,  $R^{b'}$ ,  $R^c$ ,  $R^{c'}$ ,  $R^d$ ,  $R^e$ ,  $R^j$ ,  $R^n$ ,  $R^{n'}$ ,  $R^p$ ,  $R^{p'}$ ,  $R^t$ ,  $R^u$ ,  $R^v$ ,  $R^{v'}$ ,  $R^e$ ,  $R^h$ ,  $R^k$ ,  $R^t$ ,  $R^f$ ,  $R^i$ ,  $R^m$  and  $R^u$  and least one such group includes a hydroxy substituent.
12. (Currently Amended) A compound according to claim 5 wherein  $R^9$  is selected from a ~~group of formula~~ (1), (3), (6) or (10).


13. (Currently Amended) A compound according to any one of ~~the preceding~~ claims 1, 4, 5 or 6, wherein X is NH or O.
14. (Canceled)
15. (Canceled)
16. (Canceled)
17. (Canceled)
18. (Currently Amended) A compound according to claim ~~137~~ wherein R<sup>5</sup> is a group of formula (iii).
19. (Canceled)
20. (Canceled)
21. (Currently Amended) A compound according to claim ~~120~~ wherein R<sup>5</sup> is substituted by a group of sub formula (II) which is a compound of formula (IIA)



where s', q' and R<sup>70</sup> are as defined in claim ~~120~~.

22. (Currently Amended) A compound according to claim ~~120~~ or claim 21 wherein ~~the substituent~~ R<sup>80</sup> includes a group R<sup>70</sup> and said group is phenyl optionally substituted by halo.

23. (Currently Amended) A compound according to claim ~~120~~ where  $R^5$  is substituted by a group of formula (d) or (e) and  $R^{100}$  is a group- $R^{70}$ -selected from optionally substituted phenyl or optionally substituted pyridyl.
24. (Currently Amended) A compound according to claim ~~120~~ or claim 23 wherein  $R^5$   $R^{80}$  is substituted by a group of sub-formula (d).
25. (Currently Amended) A compound according to any one of the preceding claims 1, 4, 5 or 6, which is a phosphate prodrug of a compound of formula (I).
26. (Currently Amended) A method for preparing a compound of formula (I) as defined in claim 1, which method comprises reacting a compound of formula (VII)



(VII)

where  $R^{1'}$ ,  $R^{2''}$ ,  $R^{3'}$ , and  $R^{4'}$  are equivalent to a group  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  as defined in relation to formula (I) or a precursor thereof, and  $R^{85}$  is a leaving group, with a compound of formula (VIII)



where X and  $R^5$  are as defined in relation to formula (I): and thereafter if desired or necessary converting a group  $R^{1'}$ ,  $R^{2''}$ ,  $R^{3'}$  or  $R^{4'}$  to a group  $R^1$ ,  $R^2$ ,  $R^3$  and/or  $R^4$  respectively or to a different such group.

27. (Original) A method for inhibiting aurora 2 kinase in a warm blooded animal, such as man, in need of such treatment, which comprises administering to said animal an

effective amount of a compound according to claim 1, or salt, ester amide or prodrug thereof.

28. (Canceled)

29. (Currently Amended) A pharmaceutical composition comprising a compound according to any one of claims 1, 4, 5 or 6, or salt, ester amide or prodrug thereof, -in combination with a pharmaceutically acceptable carrier.

30. (Cancelled)